

An Elementary Proof of the Fundamental Theorem of Tropical Algebra

Nathan Grigg*
Nathan Manwaring

February 5, 2008

Abstract

In this paper we give an elementary proof of the Fundamental Theorem of Algebra for polynomials over the rational tropical semi-ring. We prove that, tropically, the rational numbers are algebraically closed. We provide a simple algorithm for factoring tropical polynomials of a single variable. A central idea is the concept of *least-coefficient polynomials* as representatives for classes of functionally equivalent polynomials. This idea has importance far beyond the proof of the Fundamental Theorem of Tropical Algebra.

1 Introduction

In this paper we will consider the tropical semi-ring, as discussed by Richter-Gebert, Sturmfels, and Theobald in [3] and by Speyer and Sturmfels in [4]. Our goal is to give an elementary proof of the Fundamental Theorem of Algebra as it applies to the tropical semi-ring.

Although the authors of some papers refer to this theorem, they do not do more than confirm it as true or dismiss it as trivial. Nevertheless, our proof of this theorem is key to understanding vital components of the tropical algebraic structure. We note that one version of the proof has been published by Izhakian in [2], but Izhakian gives his proof over an “extended” tropical semi-ring that is substantially different from the standard tropical semi-ring that most others study. Hence, there is merit in discussing this elementary proof and the underlying ideas it addresses.

Fundamental Theorem of Tropical Algebra. *Every tropical polynomial in one variable with rational coefficients can be factored uniquely as a product of linear tropical polynomials with rational coefficients, up to functional equivalence.*

*Nathan Grigg and Nathan Manwaring are undergraduates at Brigham Young University. This research was funded in part by a grant from the BYU Office of Research and Creative Activities.

It is important to note that this theorem only applies up to functional equivalence. To illustrate this, note that we would factor $x^2 \oplus 4x \oplus 6$ as $(x \oplus 3)^2$. As functions, these are the same—for any x they are equal. Nevertheless, the second expression expands to the polynomial $x^2 \oplus 3x \oplus 6$, which is not the same polynomial as the first. For this reason, together with the fact that geometric properties of a polynomial depend only on its function, we will regard two polynomials as equivalent if they define the same function. For more information, see Section 2.

Since we will be dealing with equivalence classes of polynomials, it is useful to have a representative for each functional equivalence class. In Section 3, we discuss one possible, very useful representative, called a *least-coefficient polynomial*. We prove that every tropical polynomial is functionally equivalent to a least-coefficient polynomial and that each least-coefficient polynomial can be easily factored using the formula given in Section 4.

Definition 1.1. The *rational tropical semi-ring* is $\mathcal{Q} = (\mathbb{Q} \cup \infty, \oplus, \odot)$, where

$$\begin{aligned} a \oplus b &:= \min(a, b), \text{ and} \\ a \odot b &:= a + b. \end{aligned}$$

We note that the additive identity of $\mathcal{Q}$ is ∞ and the multiplicative identity is 0 . Elements of $\mathcal{Q}$ do not have additive inverses, but the multiplicative inverse of a is the classical *negative* a . The commutative, associative, and distributive properties hold.

Notation We will write tropical multiplication $a \odot b$ as ab , and repeated multiplication $a \odot a$ as a^2 . We will write classical addition, subtraction, multiplication, and division as $a + b$, $a - b$, $a \cdot b$, and $\frac{a}{b}$, respectively.

2 Equality and Functional Equivalence

A polynomial $f(x) \in \mathcal{Q}[x]$ is defined to be a formal sum

$$f(x) = a_n x^n \oplus a_{n-1} x^{n-1} \oplus \cdots \oplus a_0.$$

For two polynomials f and g , we write $f = g$ if each pair of corresponding coefficients of f and g are equal.

We can also think of a tropical polynomial as a function. Two polynomials are *functionally equivalent* if for each $x \in \mathcal{Q}$, $f(x) = g(x)$. In this case, we write $f \sim g$. Notice that functional equivalence does not imply equality. For example, the polynomials $x^2 \oplus 1x \oplus 2$ and $x^2 \oplus 2x \oplus 2$ are functionally equivalent, but not equal as polynomials. In general, functional equivalence is a more useful equivalence relation to use with tropical polynomials than equality of coefficients.

Definition 2.1. A coefficient a_i of a polynomial $f(x)$ is a *least coefficient* if for any $b \in \mathcal{Q}$ with $b < a_i$, the polynomial $g(x)$ formed by replacing a_i with b is not functionally equivalent to $f(x)$.

Note. If $f(x) = a_n x^n \oplus a_{n-1} x^{n-1} \oplus \cdots \oplus a_r x^r$, where $a_n, a_r \neq \infty$, then a_n and a_r are least coefficients. Additionally, if $r < i < n$ and $a_i = \infty$, then a_i is not a least coefficient.

Lemma 2.2 (Alternate definition of least coefficient). *Let $a_i x^i$ be a term of a polynomial $f(x)$, with a_i not equal to infinity. Then a_i is a least coefficient of $f(x)$ if and only if there is some $x_0 \in \mathbb{Q}$ such that $f(x_0) = a_i x_0^i$.*

Proof. For all $x \in \mathbb{Q}$, note that $f(x) \leq a_i x^i$. Suppose that there is no x such that $f(x) = a_i x^i$. Then $f(x) < a_i x^i$ for all x . Now, let $\varphi(x) = f(x) - (i \cdot x + a_i)$. Note that φ is a piecewise-linear, continuous function that is linear over a finite number of intervals. Thus, there is an interval large enough to contain all the pieces of φ . By applying the extreme value theorem to this interval, we see that $\sup \varphi \in \varphi(\mathbb{R})$, and hence $\sup \varphi < 0$. Let $\epsilon = |\sup \varphi|$ and $b \in \mathbb{Q}$ be such that $a_i - \epsilon < b < a_i$. Then

$$f(x) - (i \cdot x + b) < f(x) - (i \cdot x + a_i) + \epsilon \leq 0$$

and therefore $f(x) < i \cdot x + b$ for all $x \in \mathbb{Q}$. Therefore, the polynomial created by replacing a_i with b is functionally equivalent to $f(x)$, so a_i is not a least coefficient.

For the other direction, suppose that there is an $x_0 \in \mathbb{Q}$ such that $f(x_0) = a_i x_0^i$. Given $b < a_i$, let $g(x)$ be $f(x)$ with a_i replaced by b . Then $g(x_0) \leq b x_0^i < a_i x_0^i = f(x_0)$, so g is not functionally equivalent to f . Therefore a_i is a least coefficient. $\square$

3 Least-coefficient polynomials

Definition 3.1. A polynomial is a *least-coefficient polynomial* if all its coefficients are least coefficients.

Lemma 3.2 (Uniqueness of least-coefficient polynomials). *Let f and g be least-coefficient polynomials. Then f is equal to g if and only if f is functionally equivalent to g .*

Proof. It is clear that $f = g$ implies $f \sim g$. For the other direction, suppose that $f \neq g$. Then for some term $a_i x^i$ of $f(x)$ and the corresponding term $b_i x^i$ of $g(x)$, we have $a_i \neq b_i$. Without loss of generality, suppose $a_i < b_i$. Since g is a least-coefficient polynomial, $g(x_0) = b_i x_0^i$ for some x_0 , by Lemma 2.2. Now,

$$f(x_0) \leq a_i x_0^i < b_i x_0^i = g(x_0),$$

so f is not functionally equivalent to g . $\square$

We will now prove that every functional equivalence class contains a unique least-coefficient polynomial. This least-coefficient representative is often the most useful way to represent a functional equivalence class of tropical polynomials.

Lemma 3.3. *Let $f(x) = a_n x^n \oplus a_{n-1} x^{n-1} \oplus \cdots \oplus a_r x^r$. There is a unique least-coefficient polynomial $g(x) = b_n x^n \oplus b_{n-1} x^{n-1} \oplus \cdots \oplus b_r x^r$ such that $f \sim g$. Furthermore, each coefficient b_j of $g(x)$ is given by*

$$b_j = \min \left(\left\{ a_j \right\} \cup \left\{ \frac{a_i \cdot (k - j) + a_k \cdot (j - i)}{k - i} \mid r \leq i < j < k \leq n \right\} \right). \quad (1)$$

Proof. First we will show that $f \sim g$. Given x_0 , note that $f(x_0) = a_s x_0^s = a_s + s \cdot x_0$ for some s . Also,

$$\begin{aligned} g(x_0) &= \min_{r \leq j \leq n} \{b_j + j \cdot x_0\} \\ &= \min_{r \leq i < j < k \leq n} \left\{ a_j + j \cdot x_0, \frac{a_i \cdot (k-j) + a_k \cdot (j-i)}{k-i} + j \cdot x_0 \right\}. \end{aligned} \quad (2)$$

So for any i, j , and k such that $r \leq i < j < k \leq n$, if $x_0 \geq \frac{a_i - a_k}{k-i}$ then

$$\begin{aligned} a_s + s \cdot x_0 &\leq a_i + i \cdot x_0 \\ &= \frac{a_i \cdot (k-j) + a_k \cdot (j-i)}{k-i} + (j-i) \cdot \left(\frac{a_i - a_k}{k-i} \right) + i \cdot x_0 \\ &\leq \frac{a_i \cdot (k-j) + a_k \cdot (j-i)}{k-i} + (j-i) \cdot x_0 + i \cdot x_0 \\ &= \frac{a_i \cdot (k-j) + a_k \cdot (j-i)}{k-i} + j \cdot x_0. \end{aligned}$$

A similar argument shows that if $x_0 \leq \frac{a_i - a_k}{k-i}$, then

$$a_s + s \cdot x_0 \leq a_k + k \cdot x_0 \leq \frac{a_i \cdot (k-j) + a_k \cdot (j-i)}{k-i} + j \cdot x_0.$$

Since this is true for all i, j , and k , the equation in (2) evaluates to $g(x_0) = a_s x_0^s$, so $g(x_0) = f(x_0)$ and $f \sim g$, as desired.

Secondly, we must show g is a least-coefficient polynomial. Given a coefficient b_j in g , suppose that a_j is a least coefficient of f . From Equation (1) we see that $b_j \leq a_j$. Since a_j is a least coefficient, there is some x_0 such that $f(x_0) = a_j x_0^j$, so $b_j x_0^j \geq g(x_0) = f(x_0) = a_j x_0^j$. Therefore $b_j = a_j$ and $g(x_0) = b_j x_0^j$.

Now suppose that a_j is not a least coefficient. Then since a_r and a_n are least-coefficient, we can choose $u < j$ and $v > j$ such that a_u and a_v are least coefficients and for any t such that $u < t < v$, a_t is not a least coefficient.

Let $x_0 = \frac{a_u - a_v}{v-u}$ and suppose, by way of contradiction, that $f(x_0) \neq a_u x_0^u$. Then $f(x_0) = a_w x_0^w < a_u x_0^u$ for some w . Note that a_w is a least coefficient, so it cannot be that $u < w < v$ by our assumption on u and v . If $w < u$ then for $x \geq x_0$,

$$\begin{aligned} a_w + w \cdot x &= a_w + u \cdot x - (u-w) \cdot x \\ &\leq a_w + u \cdot x - (u-w) \cdot x_0 \\ &= a_w + w \cdot x_0 + u \cdot (x - x_0) \\ &< a_u + u \cdot x_0 + u \cdot (x - x_0) \\ &= a_u + u \cdot x \end{aligned}$$

For $x < x_0$,

$$\begin{aligned}
a_v + v \cdot x &= a_v + u \cdot x + (v - u) \cdot x \\
&< a_v + u \cdot x + (v - u) \cdot x_0 \\
&= a_v + v \cdot x_0 + u \cdot (x - x_0) \\
&= a_u + u \cdot x_0 + u \cdot (x - x_0) \\
&= a_u + u \cdot x
\end{aligned}$$

So there is no x such that $f(x) = a_u x^u$ and thus a_u is not a least coefficient, which contradicts our assumption. If $w > v$, a similar argument shows that a_v is not a least coefficient, again contradicting our assumption. Therefore,

$$\begin{aligned}
f(x_0) &= a_u + u \cdot \left(\frac{a_u - a_v}{v - u} \right) \\
&= \frac{a_u \cdot (v - j) + a_v \cdot (j - u)}{v - u} + j \cdot \left(\frac{a_u - a_v}{v - u} \right) \\
&= c + j \cdot x_0, \text{ where } c = \frac{a_u \cdot (v - j) + a_v \cdot (j - u)}{v - u}
\end{aligned} \tag{3}$$

Again, from (1) we see that $b_j \leq c$, and from (3) we see $cx_0^j = f(x_0) = g(x_0) \leq b_j x_0^j$. So $b_j = c$ and $g(x_0) = b_j x_0^j$.

Finally, g is the only such polynomial by Lemma 3.2. $\square$

Note. The use of a least-coefficient polynomial as a best representative for a functional equivalence class is one of the key ideas of this paper. We cannot develop well-defined algebraic transformations of tropical polynomials without unique representatives for functional equivalence classes. While Izhakian discusses in [2] what he calls an “effective” coefficient (similar to a least coefficient), the idea of using least-coefficient polynomials to represent functional equivalence classes has not been discussed.

Lemma 3.4. *Let $f(x) = a_n x^n \oplus a_{n-1} x^{n-1} \oplus \dots \oplus a_r x^r$, where each a_i is not infinity. Let $d_i = a_{i-1} - a_i$ be the difference between two consecutive coefficients. Then $f(x)$ is a least-coefficient polynomial if and only if the difference between consecutive coefficients is non-decreasing, that is, if $d_n \leq d_{n-1} \leq \dots \leq d_{r+1}$.*

Proof. Suppose that f has a set of consecutive coefficients whose differences are decreasing, that is, ax^{i+1} , bx^i , and cx^{i-1} are consecutive terms of $f(x)$ such that $b - a > c - b$. Then $b > \frac{1}{2} \cdot (a + c)$. We will show that $f(x_0) < bx_0^i$ for all x_0 , meaning that b is not a least coefficient.

Given x_0 , if $x_0 \leq \frac{1}{2} \cdot (c - a)$ then

$$\begin{aligned}
ax_0^{i+1} &= (i+1) \cdot x_0 + a \\
&\leq i \cdot x_0 + \frac{1}{2} \cdot (c - a) + a \\
&= i \cdot x_0 + \frac{1}{2} \cdot (c + a) \\
&< i \cdot x_0 + b = bx_0^i,
\end{aligned}$$

so $f(x_0) \leq ax_0^{i+1} < bx_0^i$. Similarly, if $x_0 \geq \frac{1}{2} \cdot (c - a)$,

$$\begin{aligned} cx_0^{i-1} &= (i-1) \cdot x_0 + c \\ &\leq i \cdot x_0 - \frac{1}{2} \cdot (c - a) + c \\ &= i \cdot x_0 + \frac{1}{2} \cdot (c + a) \\ &< i \cdot x_0 + b = bx_0^i, \end{aligned}$$

so $f(x_0) \leq cx_0^{i-1} < bx_0^i$. Therefore b is not a least coefficient, and f is not a least-coefficient polynomial.

For the other direction, suppose that the differences between the coefficients of $f(x)$ are nondecreasing. Since $a_n, a_r \neq \infty$, a_n and a_r are least coefficients. Let a_i be a coefficient of f , with $r < i < n$, and let $x_0 = \frac{a_{i-1} - a_{i+1}}{2}$. We will show that $f(x_0) = a_i x_0^i$, so a_i is a least coefficient. We must show for all k that $i \cdot x_0 + a_i \leq k \cdot x_0 + a_k$.

This is certainly true for $i = k$. Suppose $k > i$. Then, since $(a_t - a_{t+1}) \leq (a_s - a_{s+1})$ for $t \geq s$, we have

$$\begin{aligned} (a_i - a_{i+2}) &= (a_i - a_{i+1}) + (a_{i+1} - a_{i+2}) \leq 2 \cdot (a_i - a_{i+1}) \\ (a_i - a_{i+3}) &= (a_i - a_{i+2}) + (a_{i+2} - a_{i+3}) \leq 3 \cdot (a_i - a_{i+1}) \end{aligned}$$

And in general we get

$$\begin{aligned} (a_i - a_k) &\leq (a_i - a_{i+1}) \cdot (k - i) = \frac{1}{2} \cdot (2 \cdot (a_i - a_{i+1})) \cdot (k - i) \\ &\leq \frac{1}{2} \cdot ((a_{i-1} - a_i) + (a_i - a_{i+1})) \cdot (k - i) = x_0 \cdot (k - i). \end{aligned}$$

Thus, $i \cdot x_i + a_i \leq k \cdot x_i + a_k$. A similar argument holds for $k < i$. So (tropically) $a_i x_i^i \leq a_s x_i^s$ for all s . This means that $f(x_i) = a_i x_i^i$, so a_i is a least coefficient. Therefore, f is a least-coefficients polynomial. $\square$

Note. If $f(x)$ has a coefficient a_i such that $a_i = \infty$ for $r < i < n$, then f is not a least-coefficient polynomial; but of course, $a_i = \infty$ for all $i > n$ and all $i < r$, even in a least-coefficient polynomial.

4 The Fundamental Theorem of Tropical Algebra

Fundamental Theorem of Tropical Algebra. *Let $f(x) = a_n x^n \oplus a_{n-1} x^{n-1} \oplus \dots \oplus a_r x^r$ be a least coefficients polynomial. Then $f(x)$ can be written uniquely as the product of linear factors*

$$a_n x^r (x \oplus d_n) (x \oplus d_{n-1}) \cdots (x \oplus d_{r+1}), \quad (4)$$

where $d_i = a_{i-1} - a_i$. In other words, the roots of $f(x)$ are the differences between consecutive coefficients.

Proof. Since $f(x)$ is a least-coefficient polynomial, the differences between consecutive coefficients is non-decreasing, i.e., $d_n \leq d_{n-1} \leq \dots \leq d_{r+1}$. Knowing these inequalities, we can expand (4) to get

$$a_n x^n \oplus a_n d_n x^{n-1} \oplus a_n d_n d_{n-1} x^{n-2} \oplus \dots \oplus a_n d_n d_{n-1} \dots d_{r+1} x. \quad (5)$$

But the coefficient of the x^i term in this polynomial is

$$a_n d_n d_{n-1} \dots d_{i+1} = a_n + d_n + d_{n-1} + \dots + d_{i+1}.$$

A straightforward computation shows that this is equal to a_i , so the polynomial in (5) is equal to $f(x)$, as desired.

Now suppose that there is another way of writing $f(x)$ as a product of linear factors. Call this product g and note that it must have the same degree as f . Additionally, the smallest non-infinite term of g must have the same degree as the smallest non-infinite term of f . Hence, we are able to write $g(x) = a'_n x^n (x \oplus d'_n) (x \oplus d'_{n-1}) \dots (x \oplus d'_{r+1})$, with each d'_i chosen, after reindexing, if necessary, such that $d'_n \leq d'_{n-1} \leq \dots \leq d'_{r+1}$. Expanding this product shows that the differences between consecutive coefficients of g are non-decreasing, so g is a least-coefficient polynomial by Lemma 3.4. We see from (5) that $f \neq g$, so by Lemma 3.2, f is not functionally equivalent to g . Therefore, the factorization is unique. $\square$

Finally we note that tropical factoring gives us a slightly different result than classical factoring. Classically, the set of roots (or zero locus) of a polynomial is the set of points at which the polynomial evaluates to the additive identity. Unfortunately, tropical polynomials have either no roots or trivial roots in this sense. In fact, if $f(x) \neq \infty$, then $f(x_0)$ never evaluates to the additive identity ∞ when $x_0 \neq \infty$. However, as we have seen, polynomials in $\mathcal{Q}[x]$ can be factored and seem to have “roots,” although they do not evaluate to the additive identity at these points. Clearly, we must use a different, more meaningful definition. In [3] motivation is given for the following definition of zero locus.

Definition 4.1. Let $f(x) \in \mathcal{Q}[x]$. The *tropical zero locus* (or *corner locus*) $\mathcal{Z}(f)$ is the set of points x_0 in $\mathcal{Q}$ for which at least two monomials of f attain the minimum value.

The d_i in (4) are precisely that points of $\mathcal{Z}(f)$, as we now show.

Theorem 4.2. Given a point $d \in \mathcal{Q}$ and a least-coefficient polynomial $f(x)$, $x \oplus d$ is a factor of $f(x)$ if and only if $f(d)$ attains its minimum on at least two monomials.

Proof. First, suppose that $x \oplus d$ is a factor of $f(x)$. If we write f as a product of linear factors as in (4), $d_i = d$ for some d_i . For all $j < i$,

$$\begin{aligned} a_i + i \cdot d_i &= a_i + (i-j) \cdot d_i + j \cdot d_i \\ &\leq a_i + \underbrace{d_i + d_{i-1} + \dots + d_{j+1}}_{i-j \text{ terms}} + j \cdot d_i \\ &= a_j + j \cdot d_i. \end{aligned}$$

A similar calculation shows that for $j > i$, we have $a_i + i \cdot d_i \leq a_j + j \cdot d_i$. So $f(d_i) = a_i d_i^i$. Also, $a_i d_i^i = a_n d_n d_{n-1} \cdots d_{i+1} d_i^i = a_n d_n d_{n-1} \cdots d_i d_i^{i-1} = a_{i-1} d_i^{i-1}$, so the minimum is attained by at least two monomials of $f(x)$ at $x = d$.

For the other direction, suppose that the minimum is attained by two monomials at $f(d)$. By way of contradiction, suppose that these monomials are not consecutive. Then for some $j < i < k$, we have $a_j d^j = a_k d^k < a_i d^i$. If $x \leq d$ then

$$\begin{aligned} a_k + k \cdot x &= a_k + i \cdot x + (k - i) \cdot x \leq a_k + i \cdot x + (k - i) \cdot d \\ &= a_k + k \cdot d + i \cdot (x - d) < a_i + i \cdot d + i \cdot (x - d) \\ &= a_i + i \cdot x \end{aligned}$$

Similarly, if $x \geq d$, then $a_j x^j < a_i x^i$. Thus there is no x such that $f(x) = a_i x^i$, so a_i is not a least coefficient, which is a contradiction. Therefore there is some i such that $a_i d^i = a_{i-1} d^{i-1}$. Thus we have

$$0 = a_{i-1} + (i - 1) \cdot d - (a_i + i \cdot d) = a_{i-1} - a_i - d.$$

So $d = a_{i-1} - a_i$, the difference between two consecutive coefficients. Since f is a least-coefficient polynomial, $x \oplus d$ is a factor of f by the Fundamental Theorem. $\square$

Thus, as in the classical case, the unique factorization of a polynomial in $\mathbb{Q}[x]$ gives us what could be considered the *roots* of the polynomial. It is clear that all of the arguments and results of this paper hold if we replace the rationals $\mathbb{Q}$ with any ordered field. Thus any ordered field, together with ∞ , can be said to be tropically algebraically closed.

References

- [1] Nathan Grigg, *Factorization of tropical polynomials in one and several variables*, Honor's Thesis, Brigham Young University, June 2007.
- [2] Zur Izhakian, *Tropical varieties, ideals and an algebraic nullstellensatz*, preprint <http://arxiv.org/pdf/math.AC/0511059>, 2005.
- [3] Jürgen Richter-Gebert, Bernd Sturmfels, and Thorsten Theobald, *First steps in tropical geometry*, Idempotent mathematics and mathematical physics, Contemp. Math., vol. 377, Amer. Math. Soc., Providence, RI, 2005, pp. 289–317. MR MR2149011 (2006d:14073)
- [4] David Speyer and Bernd Sturmfels, *Tropical mathematics*, preprint <http://arxiv.org/pdf/math.CO/0408099>, 2004.